

U-CyTech B.V.
Yalelaan 48
3584 CM Utrecht
The Netherlands
+31 85 073 1460
info@ucytech.com
www.ucytech.com

Data sheet

Monoclonal antibody to monkey interleukin 13 (IL-13) FITC-labeled

Cat. No.	CT120
Clone	QS-13
Isotype	Mouse IgG ₁ , κ
Immunogen	Recombinant human IL-13
Production	<i>In vitro</i> using serum free medium
Purification	Ion exchange chromatography
Labeling	With 5(6)-carboxyfluorescein-N-hydroxysuccinimide ester (FLUOS)
Packaging	Lyophilized and vacuum-packed
Contents	50 µg/vial
Buffer	Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose.
Specificity:	Natural IL-13 of rhesus macaques, cynomolgus monkeys, pig-tailed macaques, Japanese macaques, barbary macaques, lion-tailed macaques, baboons, African green monkeys and marmosets binds with high efficiency.
Sterility	Membrane filtered (0.2 µm)
Reconstitution	Dissolve the contents of the vial by injection of 0.5 ml sterile distilled water.
Working conc.	1 to 2 µl per 10 ⁶ PBMC (minimal 250 tests)
Stability	Lyophilized product is stable for at least one year at 4°C (expiry date is indicated on the vial). After reconstitution, the contents can be safely stored at 4°C for one month or for one year at -20°C. Add 0.02% sodium azide to prevent bacterial growth.
Quantitation	Antibody concentration was determined by absorbance, taking A ₂₈₀ =1.4 for a 1 mg/ml solution.
Application	Flow cytometry

References can be found in the Reference Database at www.ucytech.com/references.

