

U-CyTech B.V.
Yalelaan 42
3584 CM Utrecht
The Netherlands
+31 85 073 1460
info@ucytech.com
www.ucytech.com

Data sheet

Monoclonal antibody to monkey CD3 FITC-labeled

Cat. No.	CT163
Clone	FN-18
Isotype	Mouse IgG ₁
Immunogen	Rhesus monkey peripheral blood mononuclear cells
Production	<i>In vitro</i> using serum free medium
Purification	Ion exchange chromatography
Labeling	With 5(6)-carboxyfluorescein-N-hydroxysuccinimide ester (FLUOS).
Packaging	Lyophilized and vacuum-packed
Contents	50 µg/vial
Buffer	Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose.
Specificity	Rhesus macaque, cynomolgus monkey and baboon
Sterility	Membrane-filtered (0.2 µm)
Reconstitution	Dissolve the contents of the vial by injection of 0.5 ml sterile distilled water. Working dilution: 10x in PBS + 1% BSA.
Stability	Lyophilized product, protected from light, is stable for ≥1 year at 2-8°C (see vial for expiry). After reconstitution, store at 2-8°C for up to one month if sterile. For long-term use, aliquot the solution, protect from light and freeze at -32 to -18°C (stable ≥1 year). Add 0.02% sodium azide to prevent bacterial growth.
Quantitation	Antibody concentration was determined by absorbance, taking A ₂₈₀ =1.4 for a 1 mg/ml solution.
Application	Flowcytometry (20 to 50 µl per 10 ⁶ cells) Fluorescence microscopy

