

U-CyTech BV

Yalelaan 48
3584 CM Utrecht
The Netherlands
P +31.30.253 5960
F +31.30.253 9344
INFO@ucytech.com
www.ucytech.com

Data sheet Marmoset TNF-α ELISA antibody pair; 20-plate format

Cat. No.: CT772-20

Coating antibodies (4 vials)

Product: Monoclonal antibody to marmoset tumor necrosis factor alpha (TNF-α)

Isotype: Mouse IgG₁

Production: In vitro using serum free medium
Purification: Ion exchange chromatography

Buffer: Prior to lyophilization: 0.25 ml PBS + 125 mM trehalose

Application: Inject 0.25 ml distilled water into the vial and dilute 100 times in PBS. The content

of one vial is sufficient for five 96-well ELISA plates (480 determinations; 50

μl/well).

Detection antibodies (4 vials)

Product: Biotinylated monoclonal antibody to marmoset tumor necrosis factor alpha (TNF-α)

Isotype: Mouse IgG₁

Production: In vitro using serum free medium
Purification: Ion exchange chromatography

Labeling: With Biotin-7-NHS (N-hydroxysuccinimide)

Buffer: Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose

Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA +

0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates

(480 determinations; 100 µl/well).

Standards (10 vials)

Product: Recombinant tumor necrosis factor alpha (TNF-α)

Application: Cytokine standard in an ELISA system

Reconstitution: Inject 0.5 ml distilled water into the vial. Use immediately.

Conjugate (4 vials)

Product: SPP conjugate (Streptavidin-HRP)

Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA +

0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates

(480 determinations; 100 μ l/well).

The product should be used in combination with TMB substrate.

General

Sensitivity: 1 pg/ml

Specificity: Validated for detecting natural marmoset TNF- α .

Standards are calibrated with natural marmoset TNF- α .

Sterility: Membrane filtered (0.2 µm)

Stability: Lyophilized SPP conjugate is stable for at least one year at -20°C in the dark, the

other lyophilized products are stable for at least one year at 4°C.

After reconstitution, the antibodies are stable for one year at $4\,^\circ\text{C}$ (if kept sterile) and SPP for minimal one year at $\text{-}20\,^\circ\text{C}$ in the dark. The reconstituted standard

preparation should be used immediately.



For research use only Version 140820