

# U-CyTech BV

Yalelaan 48  
3584 CM Utrecht  
The Netherlands  
P +31.30.253 5960  
F +31.30.253 9344  
INFO@ucytech.com  
www.ucytech.com

## Data sheet Human IL-17A ELISA antibody pair; 20-plate format

Cat. No.: CT566-20

### Coating antibodies (4 vials)

Product: Monoclonal antibody to human interleukin 17A (IL-17A)  
Isotype: Mouse IgG<sub>2b</sub>  
Production: *In vitro* using serum free medium  
Purification: Ion exchange chromatography  
Buffer: Prior to lyophilization: 0.25 ml PBS + 125 mM trehalose  
Application: Inject 0.25 ml distilled water into the vial and dilute 100 times in PBS. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 50 µl/well).

### Detection antibodies (4 vials)

Product: Biotinylated monoclonal antibody to human interleukin 17A (IL-17A)  
Isotype: Mouse IgG<sub>2b</sub>  
Production: *In vitro* using serum free medium  
Purification: Protein A-affinity chromatography  
Labeling: With Biotin-7-NHS (N-hydroxysuccinimide)  
Buffer: Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose  
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 µl/well).

### Standards (10 vials)

Product: Recombinant human interleukin 17A (IL-17A)  
Application: Cytokine standard in an ELISA system  
Reconstitution: Dissolve the contents of one vial by injection of 0.5 ml distilled water into the vial. Use immediately.

### Conjugate (4 vials)

Product: SPP conjugate (Streptavidin-HRP)  
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 µl/well).  
The product should be used in combination with TMB substrate.

### General

Sensitivity: 2 pg/ml  
Specificity: Validated for detecting natural and recombinant human IL-17A  
Sterility: Membrane filtered (0.2 µm)  
Stability: Lyophilized SPP conjugate is stable for at least one year at -20°C in the dark, the other lyophilized products are stable for more than one year at 4°C. After reconstitution, the antibodies are stable for one year at 4°C (if kept sterile) and SPP for minimal one year at -20°C in the dark. The reconstituted standard preparation should be used immediately.