

U-CyTech BV

Yalelaan 48
3584 CM Utrecht
The Netherlands
P +31.30.253 5960
F +31.30.253 9344
INFO@ucytech.com
www.ucytech.com

Data sheet Mouse IFN- γ ELISA antibody pair; 10-plate format

Cat. No.: CT755-10

Coating antibodies (2 vials)

Product: Monoclonal antibody to mouse interferon gamma (IFN- γ)
Isotype: Rat IgG_{2a}
Production: *In vitro* using serum free medium
Purification: Ammonium sulphate precipitation and DEAE ion-exchange chromatography
Buffer: Prior to lyophilization: 0.25 ml PBS + 125 mM trehalose
Application: Inject 0.25 ml distilled water into the vial and dilute 100 times in PBS. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 80 μ l/well).

Detection antibodies (2 vials)

Product: Biotinylated polyclonal antibody to mouse interferon gamma (IFN- γ)
Isotype: Rabbit IgG
Purification: Ammonium sulphate precipitation and protein A- and ligand-affinity chromatography
Labeling: With Biotin-7-NHS (N-hydroxysuccinimide)
Buffer: Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 μ l/well).

Standards (5 vials)

Product: CHO-derived recombinant mouse interferon gamma (IFN- γ)
Application: Cytokine standard in an ELISA system
Reconstitution: Inject 0.5 ml distilled water into the vial. Use immediately.

Conjugate (2 vials)

Product: SPP conjugate (Streptavidin-HRP)
Application: Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA + 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plates (480 determinations; 100 μ l/well).
The product should be used in combination with TMB substrate.

General

Sensitivity: 1 pg/ml
Specificity: Validated for detecting natural and recombinant mouse IFN- γ .
Sterility: Membrane filtered (0.2 μ m)
Stability: Lyophilized SPP conjugate is stable for at least one year at -20°C in the dark, the other lyophilized products are stable for at least one year at 4°C.
After reconstitution, the antibodies are stable for one year at 4°C (if kept sterile) and SPP for minimal one year at -20°C in the dark. The reconstituted standard preparation should be used immediately.
References: Geluk, A. *et al.* 2000. J. Immunol. 165: 6463-6471