

Data sheet Mouse IFN- γ ELISA antibody pair; 20-plate format

U-CyTech BV Yalelaan 48 3584 CM Utrecht The Netherlands P +31.30.253 5960 F +31.30.253 9344 INFO@ucytech.com

Cat. No.:	CT755-20
Coating antibo	dies (4 vials)
Product:	Monoclonal antibody to mouse interferon gamma (IFN-y)
lsotype:	Rat IgG _{2a}
Production:	In vitro using serum free medium
Purification:	Ammonium sulphate precipitation and DEAE ion-exchange chromatography
Buffer:	Prior to lyophilization: 0.25 ml PBS + 125 mM trehalose
Application:	Inject 0.25 ml distilled water into the vial and dilute 100 times in PBS. The cont of one vial is sufficient for five 96-well ELISA plates (480 determinations; 50 μ l/well).
Detection antil	bodies (4 vials)
Product:	Biotinylated polyclonal antibody to mouse interferon gamma (IFN- γ)
lsotype:	Rabbit IgG
Purification:	Ammonium sulphate precipitation and protein A- and ligand-affinity chromatography
Labeling:	With Biotin-7-NHS (N-hydroxysuccinimide)
Buffer:	Prior to lyophilization: 0.5 ml PBS + 1% BSA + 125 mM trehalose
Application:	Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plate (480 determinations; 100 μ l/well).
Standards (10	vials)
Product:	CHO-derived recombinant mouse interferon gamma (IFN- γ)
Application:	Cytokine standard in an ELISA system
Reconstitution:	Inject 0.5 ml distilled water into the vial. Use immediately.
Conjugate (4 v	ials)
Product:	SPP conjugate (Streptavidin-HRP)
Application:	Inject 0.5 ml distilled water into the vial and dilute 100 times in PBS + 0.5% BSA 0.05% Tween-20. The content of one vial is sufficient for five 96-well ELISA plate (480 determinations; 100 μ l/well). The product should be used in combination with TMB substrate.
General	
Sensitivity:	1 pg/ml
Specificity:	Validated for detecting natural and recombinant mouse IFN-y.
Sterility:	Membrane filtered (0.2 µm)
Stability:	Lyophilized SPP conjugate is stable for at least one year at -20° C in the dark, th other lyophilized products are stable for at least one year at 4° C. After reconstitution, the antibodies are stable for one year at 4° C (if kept steril and SPP for minimal one year at -20° C in the dark. The reconstituted standard preparation should be used immediately.

